arXiv:1108.1467v2 [math.FA] 14 Nov 2011

KY FAN INEQUALITIES

MOHAMMAD SAL MOSLEHIAN

Dedicated to the memory of Professor Ky Fan

Abstract. There are several inequalities in the literature carrying the name of Ky Fan. We survey these well-known Ky Fan inequalities and some other significant inequalities generalized by Ky Fan and review some of their recent developments.

1. Introduction

According to MathSciNet, Professor Ky Fan (1914-2010) published 126 papers and books and coauthored with 15 mathematicians. His earliest indexed item goes back to 1940 [19]. It is notable that contributions of Ky Fan to mathematics have provided a lot of influence in the development of nonlinear analysis, convex analysis, approximation theory, operator theory, linear algebra, mathematical programming and mathematical economics; see e.g. [55]. In the literature, there are several inequalities due to Ky Fan in various fields; cf. [12]. In this article we try to briefly survey most important ones by using some information in MathSciNet and Zentralblatt MATH for some unavailable old papers.

2. Preliminaries on matrix analysis

An n-tuple $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ with $x_1 \geq \cdots \geq x_n$ is said to be weakly majorized by an n-tuple $y = (y_1, \dots, y_n)$ of real numbers with $y_1 \geq \cdots \geq y_n$, denoted by $x \prec_w y$, if $\sum_{i=1}^k x_i \leq \sum_{i=1}^k y_i$ for $k = 1, \dots, n$. If, in addition,

2010 *Mathematics Subject Classification.* Primary 00-02; Secondary 26D15, 47A63.

Key words and phrases. Ky Fan inequalities, eigenvalue inequality, matrix inequality, minimax inequality, M-Matrix, Szász's inequality, Ky Fan norm, Kantorovich inequality, Ky Fan–Todd determinantal inequality, Ky Fan–Taussky–Todd inequality.

$\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$, then x is said to be majorized by y and it is denoted by $x \prec y$. Let $\mathrm{Re}(x)$ stand for $(\mathrm{Re}(x_1), \ldots, \mathrm{Re}(x_n))$ for an $x = (x_1, \ldots, x_n) \in \mathbb{C}^n$.

Let $\mathbb{M}_n$ be the matrix algebra of all $n \times n$ matrices with entries in the complex field identified with the algebra $\mathbb{B}(\mathbb{C}^n)$ of all linear operators on the Hilbert space $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$. We denote by $\mathbb{H}_n$ the set of all Hermitian matrices in $\mathbb{M}_n$. By I_n (or I if there is no ambiguity) we denote the identity matrix of $\mathbb{M}_n$. A matrix $A \in \mathbb{H}_n$ is called positive-semidefinite if $\langle Ax, x \rangle \geq 0$ holds for every $x \in \mathbb{C}^n$ and then we write $A \geq 0$. In particular, if A is invertible, then it is called positive-definite and we write $A > 0$. The absolute value of A is defined by $|A| = (A^*A)^{1/2}$, where A^* denotes the conjugate transpose of A. For $A, B \in \mathbb{H}_n$, we say $A \leq B$ if $B - A \geq 0$.

For a matrix $A \in \mathbb{H}_n$, we denote by $\lambda_1(A) \geq \lambda_2(A) \geq \cdots \geq \lambda_n(A)$ the eigenvalues of A arranged in the decreasing order with their multiplicities counted. The notation $\lambda(A)$ stands for the row vector $(\lambda_1(A), \lambda_2(A), \ldots, \lambda_n(A))$. The eigenvalue inequality $\lambda(A) \leq \lambda(B)$ means $\lambda_j(A) \leq \lambda_j(B)$ $(j = 1, 2, \ldots, n)$. The trace and determinant of a matrix $A \in \mathbb{M}_n$ are defined by $\mathrm{tr}(A) = \sum_{i=1}^n \lambda_i(A)$ and $\det(A) = \prod_{i=1}^n \lambda_i(A)$, respectively. A norm $|||\cdot|||$ on $\mathbb{M}_n$ is said to be unitarily invariant if $|||UAV||| = |||A|||$ for all $A \in \mathbb{M}_n$ and all unitary matrices $U, V \in \mathbb{M}_n$. The Ky Fan norms, defined as $\|A\|_{(k)} = \sum_{j=1}^k s_j(A)$ for $k = 1, 2, \ldots, n$, provide a significant family of unitarily invariant norms; see also [11, 56]. Here $s(A) = (s_1(A), \ldots, s_n(A))$ denotes the n-tuple of the singular values of A, i.e. the eigenvalues of $|A|$, arranged in the decreasing order.

3. Ky Fan matrix inequalities

The contributions of Ky Fan to matrix theory and operator theory are substantial. In this section we study matrix inequalities due to Ky Fan. The interested reader is referred to [56, 11, 50] to see details of proofs.

Ky Fan [20, 1949] proved that if $H, K \in \mathbb{H}_n$, then

$$\lambda(H + K) \prec \lambda(H) + \lambda(K) \qquad \text{(Ky Fan eigenvalue inequality)},$$

see also [69, Theorme 7.14]. An extension of this result was given by Aujla and Silva [8] as $\lambda(f(\alpha A+(1-\alpha)B)) \prec_w \lambda(\alpha f(A)+(1-\alpha)f(B))$, where f is a convex function on an interval J and $A, B \in \mathbb{H}_n$ with spectra in J. In addition, Ky Fan [20, 1949] showed that if $A \in \mathbb{M}_n$, t is a positive integer and the eigenvalues $\kappa_i^{(t)}$ of $(A^t)^* A^t$ are arranged in the decreasing order $\kappa_1^{(t)} \geq \cdots \geq \kappa_n^{(t)}$, then

$$\kappa_1^{(t)} + \cdots + \kappa_q^{(t)} \leq \kappa_1^t + \cdots + \kappa_q^t \quad (1 \leq q \leq n)\,,$$

where $k_1, \ldots, k_n$ are eigenvalues of A^*A. Ky Fan [21, 1950] established two results which are powerful tools in obtaining matrix inequalities. They assert that if $H \in \mathbb{H}_n$, then

$$\max_{UU^*=I_k} \operatorname{tr}(UHU^*) = \sum_{i=1}^{k} \lambda_i(H), \quad \min_{UU^*=I_k} \operatorname{tr}(UHU^*) = \sum_{i=1}^{k} \lambda_{n-i+1}(H) \quad (1 \leq k \leq n)$$

where maximum and minimum are taken over all $k \times n$ complex matrices U satisfying $UU^* = I_k$.

The Ky Fan dominance theorem [22, 1951] states that for $A, B \in \mathbb{M}_n$, the inequalities $\|A\|_{(k)} \leq \|B\|_{(k)}$ $(1 \leq k \leq n)$ hold if and only if $|||A||| \leq |||B|||$ for all unitarily invariant norms $|||\cdot|||$; see [57] for an application of the Ky Fan dominance theorem.

Ky Fan [22, 1951] extended an inequality of Weyl [67] (the case $m = 1$) and von Neumann [65] (the case $m = 2$): Let $A_1, \ldots, A_m \in \mathbb{M}_n$ and let $s_{j1} \geq s_{j2} \geq \cdots \geq s_{jn}$ be the singular values of A_j. Then

$$\max \left| \sum_{i=1}^{n} \langle U_1 A_1 \cdots U_m A_m x_i, x_i \rangle \right| \leq \sum_{i=1}^{n} (s_{1i} \cdots s_{mi})$$

and

$$\max \left| \det_{1 \leq i,k \leq n} \left[\langle U_1 A_1 \cdots U_m A_m x_i, x_k \rangle \right] \right| \leq \prod_{i=1}^{n} (s_{1i} \cdots s_{mi})\,,$$

where $U_1, \ldots, U_m$ run over all unitary matrices and $\{x_1, \ldots, x_n\}$ runs over all orthonormal sets in $\mathbb{C}^n$.

Furthermore, Ky Fan showed that for any matrices $A, B \in \mathbb{M}_n$,

$$s(A + B) \prec_w s(A) + s(B) \qquad \text{(Ky Fan singular value inequality)},$$

as well as

$$s_{r+t+1}(A+B) \le s_{r+1}(A) + s_{t+1}(B),$$

where $t \ge 0, r \ge 0, r+t+1 \le n$.

In the same paper [22, 1951], it was compared by Ky Fan that the real parts and modules of the diagonal elements $a_1, \ldots, a_n$ of an arbitrary matrix $A \in \mathbb{M}_n$ by establishing that

$$\mathrm{Re}(a_1, \ldots, a_n) \prec_w (|a_1|, \ldots, |a_n|) \prec_w s(A)\,.$$

Due to the fact that any matrix is unitarily equivalent to an upper triangular matrix, one gets another Ky Fan inequality [21, 1950] for eigenvalues as

$$\mathrm{Re}(\lambda(A)) \prec \lambda(\mathrm{Re}(A)),$$

where $\mathrm{Re}(A) = (A + A^*)/2$; cf. [68, Lemma 4.20]. A similar result for the eigenvalues of $\mathrm{Im}(A) = (A - A^*)/(2\mathrm{i})$ is presented by Amir-Moez and Horn [6]. The authors of [42, 1955] proved that

$$|\lambda_i(\mathrm{Re}(A))| \le s_i(A) \qquad (i = 1, \ldots, n)$$

for any matrix $A \in \mathbb{M}_n$.

Several inequalities for eigenvalues and singular values of two Hermitian matrices H, K and $H + \mathrm{i}K$ is surveyed by Cheng, Horn and Li [14]. There are still other inequalities due to Ky Fan.

Ky Fan [24, 1954] compared the characteristic roots of diagonal blocks of a Hermitian matrix and those of the matrix itself. Ky Fan proved that if $H = \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix}$ is an $n \times n$ Hermitian block matrix and $C(H) = \begin{pmatrix} H_{11} & 0 \\ 0 & H_{22} \end{pmatrix}$ is its pinching, then $\lambda(C(H)) \prec \lambda(H)$. This is true for any pinching of H; see [11, page 50]. In the next year, Ky Fan [25, 1955] proved that for H positive-definite

$$\prod_{i=1}^{m} \lambda_{n-i+1}(H) \le \prod_{i=1}^{m} \lambda_{k-i+1}(H_{11}) \ (1 \le m \le k),$$

where the matrix H_{11} and H_{22} are $k \times k$ and $m \times m$ matrices, respectively.

Ky Fan collaborated with Hoffman [42, 1955] and proved the following: Let $|||\cdot|||$ be a unitarily invariant norm on $\mathbb{M}_n$. Then

(i) If $A = U|A|$ is the polar decomposition of A, where U is unitary, and V is any unitary matrix, then

$$|||A - U||| \le |||A - V||| \le |||A + U|||\,.$$

(ii) Let $H \in \mathbb{H}_n$. It follows from the identity $A - \mathrm{Re}(A) = \frac{A-H}{2\mathrm{i}} - \frac{(A-H)^*}{2\mathrm{i}}$ that

$$|||A - \mathrm{Re}(A)||| \le |||A - H|||\,.$$

(iii) If $H, K \in \mathbb{H}_n$ and $U = (H - \mathrm{i}I)(H + \mathrm{i}I)^{-1}$, $V = (K - \mathrm{i}I)(K + \mathrm{i}I)^{-1}$ are their Cayley transforms, then

$$|||U - V||| \le 2|||H - K|||\,.$$

Ky Fan also worked on systems of inequalities. For instance, Ky Fan [10, 1963] gave some necessary and sufficient conditions for the existence of Hermitian matrices X_j satisfying the system of linear inequalities

$$\sum_{j=1}^{n}(A_{ij}X_j + X_j{A_{ij}}^*) \ge B_i \ (1 \le i \le m) \quad \text{and} \quad \mathrm{tr}\left(\sum_{j=1}^{n} C_j X_j\right) \ge c\,,$$

where $A_{ij} \in \mathbb{M}_n$ and $B_i, C_j \in \mathbb{H}_n$ and c is a real number.

4. Results of Ky Fan involving integrals

For real functions f and g on $[0,1]$, we say that $f \prec g$ whenever $\int_0^x f(t)dt \le \int_0^x g(t)dt$ for all $0 \le x \le 1$ and $\int_0^1 f(t)dt = \int_0^1 g(t)dt$. In [43, 1954], Ky Fan and Lorentz gave sufficient and necessary conditions in terms of mixed second differences of a suitable differentiable function Φ in order that

$$\int_0^1 \Phi(t, f_1(t), \ldots, f_n(t))dt \le \int_0^1 \Phi(t, g_1(t), \ldots, g_n(t))dt$$

for any bounded functions f_i, g_i with $f_i \prec g_i$.

5. Ky Fan–Taussky–Todd inequality

The Wirtinger inequality reads as follows: If f is a periodic function of the period 2π with $f, f' \in L^2([0,2\pi])$ and $\int_0^{2\pi} f = 0$, then

$$\int_0^{2\pi} f^2 \le \int_0^{2\pi} f'^2$$

with equality if and only if $f(x) = a\cos x + b\sin x$ for some constants a, b. Ky Fan, Taussky and Todd [44, 1955] proved discrete analogs of Wirtinger type. They showed that if $a_1, \dots, a_n$ are real numbers and $a_0 = a_{n+1} = 0$, then

$$2\left(1 - \cos\frac{\pi}{n+1}\right)\sum_{k=1}^{n} a_k^2 \le \sum_{k=1}^{n+1}(a_k - a_{k-1})^2$$

with equality if and only if $a_k = c\sin(k\pi/(n+1))$ $(k = 1, \dots, n)$ for some real constant c.

They also proved that if $a_1, \dots, a_n$ are real numbers and $a_0 = 0$, then

$$2\left(1 - \cos\frac{\pi}{2n+1}\right)\sum_{k=1}^{n} a_k^2 \le \sum_{k=1}^{n}(a_k - a_{k-1})^2$$

with equality if and only if $a_k = c\sin(k\pi/(2n+1))$ $(k = 1, \dots, n)$, where c is a real constant. Both constants $2\left(1 - \cos\frac{\pi}{n+1}\right)$ and $2\left(1 - \cos\frac{\pi}{2n+1}\right)$ are best possible. The converses of these two inequalities are given by Alzer [1]. Another related inequality is that of Ozeki; see [57] for its operator versions.

6. Ky Fan–Todd determinantal inequality

Ky Fan and Todd [45, 1955] showed that if $\begin{pmatrix} a_1 & \dots & a_n \\ b_1 & \dots & b_n \end{pmatrix}$ is a matrix none of whose two-column minor is singular and (p_{ij}) is a symmetric matrix such that $p = \sum_{1\le i<j\le n} p_{ij} \ne 0$, then

$$\left[\sum_{i=1}^{n} a_i^2\right]\left[\sum_{i=1}^{n} a_i^2 \sum_{i=1}^{n} b_i^2 - \Big(\sum_{i=1}^{n} a_i b_i\Big)^2\right]^{-1} \le p^{-2}\sum_{i=1}^{n}\left[\sum_{j=1, j\ne i}^{n} p_{ij}a_j(a_jb_i - a_ib_j)^{-1}\right]^2.$$

The special case where $p_{ij} = 1$, $a_i = \cos\theta_i$ and $b_i = \sin\theta_i$ for some θ_i and all i, j was obtained by Chassan [13] by some statistical arguments. Also a refinement of it was given by Chong [15].

It is worthy noting that there are a variety of other determinantal inequalities due to Ky Fan. Ky Fan [23, 1953] (see [58, p. 214]) proved that if $|C|_k$ denotes the product of the first k-th smallest eigenvalues of a real positive-definite matrix C and if $A, B \in \mathbb{M}_n$ are real positive-definite matrices and $0 \leq \lambda \leq 1$, then

$$|\lambda A + (1-\lambda)B|_k \geq |A|_k^{\lambda} |B|_k^{1-\lambda}.$$

Also Ky Fan [26, 1955] (see [56, p. 687]) proved that if $H = \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix}$ and $K = \begin{pmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{pmatrix}$ are positive-definite $n \times n$ matrices and H_{11} and K_{11} are $k \times k$ matrices, then

$$\left(\frac{\det(H+K)}{\det(H_{11}+K_{11})}\right)^{\frac{1}{n-k}} \geq \left(\frac{\det(H)}{\det(H_{11})}\right)^{\frac{1}{n-k}} + \left(\frac{\det(K)}{\det(K_{11})}\right)^{\frac{1}{n-k}}.$$

In fact, Ky Fan studied many convex/increasing functions on certain subsets of Hermitian matrices, e.g. $f_k(H) = \sum_{i=1}^{k} \lambda_i(H)$ $(H \in \mathbb{H}_n)$, from which one can get some inequalities such as the latter one.

In the same paper [26, 1955], Ky Fan proved that if $(i_1, i_2, \ldots, i_k)$ denotes the principal submatrix of a positive-definite matrix $H \in \mathbb{H}_n$ with eigenvalues $\lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n$, then

$$\lambda_1 \lambda_2 \ldots \lambda_{h+k} \leq \det(p+1, p+2, \ldots, p+k) \prod_{i=1}^{h} \frac{\det(i, p+1, p+2, \ldots, n)}{\det(p+1, p+2, \ldots, n)}$$

in which $1 \leq h \leq p < n$ and $0 \leq k \leq n-p$ and for $k = 0$ we assume that $\det(p+1, \ldots, p+0) = 1$.

If $A \in \mathbb{M}_n$ is a real matrix such that $A + A^* > 0$, then Ky Fan [33, 1973] showed that

$$\det(\mathrm{Re}(A)) \leq \det(A) \frac{1 + \mathrm{Re}(\lambda_i(A^{-1}A^*))}{2}$$

for all i. If $n \geq 4$ and $A - A^*$ is invertible, then

$$0 < \det\left(\frac{A-A^*}{2}\right) < \det(A) \frac{1 - \mathrm{Re}(\lambda_i(A^{-1}A^*))}{2}$$

for all i.

Ky Fan [37, 1983] used the Jensen inequality to prove that if $A = H + iK$ with H and K Hermitian is normalizable in the sense that there is an invertible matrix T such that T^*AT is normal, and if $p \geq 2/n$, where n is the dimension of the Hilbert space, then $|\det(A)|^p \geq |\det(H)|^p + |\det(K)|^p$. Ky Fan showed that equality holds for $p > 2/n$ if and only if either $\det(A) = 0$ or $H = 0$ (or $k = 0$). Ky Fan [34, 1974] already proved the same result for a strictly dissipative matrix, that is, a matrix whose imaginary part $\mathrm{Im}(A) = (A - A^*)/2\mathrm{i}$ is positive-definite.

7. Ky Fan mean inequality

Suppose that $x_1, \ldots, x_n \in (0, \frac{1}{2})$, $\mathcal{A}_n := (1/n)\sum_{i=1}^n x_i, \mathcal{G}_n := (\prod_{i=1}^n x_i)^{1/n}, \mathcal{H}_n := n/\sum_{i=1}^n (1/x_i)$ and $\mathcal{A}'_n, \mathcal{G}'_n, \mathcal{H}'_n$ denote the unweighted arithmetic, geometric and harmonic means of the numbers x_i and $1-x_i$, respectively. The Ky Fan inequality for means states that

$$\mathcal{G}_n/\mathcal{G}'_n \leq \mathcal{A}_n/\mathcal{A}'_n \qquad \text{(Ky Fan mean inequality)}$$

with equality only if $x_1 = \cdots = x_n$. It first appeared in the book "Inequalities" by Beckenbach and Bellman [9]. It is proved by a forward and backward induction as used for establishing the arithmetic-geometric mean inequality. It can be also proved by applying Henrici inequality $\sum_{i=1}^n 1/(1+b_i) \geq n/(1+(\prod_{i=1}^n b_i)^{1/n})$ ($b_i \geq 1$) to $b_i = 1/x_i - 1$ ($i = 1, \ldots, n$). Utilizing the monotonicity of $(1 - x^n)/n$ as a function of n, Rooin [63] established some Ky Fan type inequalities. Applying the convexity of the function $f(x) = (e^x - \alpha)/(1 + e^x)$ ($\alpha > 0$) and the Jensen inequality, the authors of [17] generalized the Ky Fan inequality. Of course, Ky Fan [22, 1951] proved that the function

$$f(x) = \frac{(\prod_{i=1}^n (1 - x_i))^{1/n}}{\sum_{i=1}^n (1 - x_i)} \cdot \frac{\sum_{i=1}^n x_i}{(\prod_{i=1}^n x_i)^{1/n}}$$

is symmetric and Schur-convex.

Many extensions, refinements as well as some counterparts have been given by many authors. Levinson [54] extended the inequality by showing that if $\varphi(u)$ has third derivative for $0 < u < 2b$, with $\varphi''(u) \geq 0$, $0 < x_i \leq b$ and $0 < p_i$,

$i = 1, 2, \ldots, n$, then

$$\frac{\sum_{i=1}^n p_i\varphi(x_i)}{\sum_{i=1}^n p_i} - \varphi\left(\frac{\sum_{i=1}^n p_i x_i}{\sum_{i=1}^n p_i}\right) \leq \frac{\sum_{i=1}^n p_i\varphi(2b - x_i)}{\sum_{i=1}^n p_i} - \varphi\left(\frac{\sum_{i=1}^n p_i(2b - x_i)}{\sum_{i=1}^n p_i}\right).$$

The Ky Fan inequality is the special case when $\varphi(u) = \log u$, $p_i = 1$ and $b = 1/2$; see also [18]. It is noticed in [52] that Ky Fan's inequality can be deduced from an inequality proved by L. Lewent in 1908. The interested reader may be referred to the interesting survey [2] for some of generalizations and refinements of the Ky Fan inequality until 1995.

In [3], the author presented the following additive version of the Ky Fan inequality:

$$\min_{1\leq i\leq n} \frac{x_i}{1 - x_i} < \frac{\mathcal{A}'_n - \mathcal{G}'_n}{\mathcal{A}_n - \mathcal{G}_n} < \max_{1\leq i\leq n} \frac{x_i}{1 - x_i}.$$

Alzer, Ruscheweyh and Salinas [5] provided the following refinement of the Ky Fan inequality:

$$\left(\frac{\mathcal{H}_n}{\mathcal{H}'_n}\right)^{n-1} \cdot \frac{\mathcal{A}_n}{\mathcal{A}'_n} \leq \left(\frac{\mathcal{G}_n}{\mathcal{G}'_n}\right)^n.$$

Some refinements of the Ky Fan inequality including means of two or more variables are obtained by Neuman and Sandor [61]. A unified approach to Sierpiński and Ky Fan inequalities by utilizing a refined Cauchy inequality and convexity of suitable functions was presented by Alzer, Ando and Nakamura [4].

8. Ky Fan inequalities involving M-matrices

A real matrix $A = (a_{ij}) \in \mathbb{M}_n$ is an M-matrix if it can be written as $A = rI - A_1$, where r is greater than the spectral radius of A_1 and A_1 has nonnegative entries. This notion was introduced by Ostrowski [62]. Ky Fan [28, 1964] gave several equivalent definitions for the notion of M-matrix. One of them is that A is an M-matrix if and only if A is nonsingular and A^{-1} has nonnegative entries. A complex matrix $B = (b_{ij})$ dominates A if $a_{ii} \leq |b_{ii}|$ for all i and $|b_{ij}| \leq |a_{ij}|$ for all $i \neq j$. If B is also an M-matrix, then we say that B proportionally dominates A whenever there exist $p_i > 0$ such that $p_i a_{ij} \leq b_{ij}$ and $a_{ij} p_j \leq b_{ij}$ for all i, j. If $\alpha \subseteq \{1, 2, \ldots, n\}$, let $A(\alpha)$ denote the determinant of the principal submatrix

formed by the rows and columns with indices contained in α, and we assume that $A(\emptyset) = 1$.

An inequality of Ostrowski [62] states that $\det(A) \leq |\det(B)|$ when B dominates A, in particular $\prod_{i=1}^n \lambda_i(A) \leq \prod_{i=1}^n h_{ii}$, when A is an M-matrix. Ky Fan [29, 1966] extended the former inequality by showing that under conditions above on the entries of A and B if $\alpha_1, \ldots, \alpha_m$ are subsets of $\{1, 2, \ldots, n\}$ such that each of the indices $1, 2, \ldots, n$ is contained in at most k of the sets α_i, then

$$\det(A)^k / \prod_{i=1}^m A(\alpha_i) \leq |\det(B)^k / \prod_{i=1}^m B(\alpha_i)|.$$

In [30, 1967] Ky Fan proved that if A and B are M-matrices of order n and a matrix $D \in \mathbb{M}_n$ dominates $A + B$ and B proportionally dominates A, then $|\det(D)|^{1/n} \geq \det(A)^{1/n} + \det(B)^{1/n}$ and

$$|\det(D)/D(\{k+1, \ldots, n\})|^{1/k} \geq [\det(A)/A(\{k+1, \ldots, n\})]^{1/k} + [\det(B)/B(\{k+1, \ldots, n\})]^{1/k}$$

for all $1 \leq k \leq n-1$.

Also, Ky Fan [27, 1960] showed that if $A, B \in \mathbb{M}_n$ are M-matrices such that $a_{ij} \leq b_{ij}$ for all i, j, and

$$\Phi(A; \alpha, \beta, \gamma) := \frac{A(\alpha \cap \beta)A(\alpha \cap \gamma)A(\beta \cap \gamma)A(\alpha \cup \beta \cup \gamma)}{A(\alpha)A(\beta)A(\gamma)A(\alpha \cap \beta \cap \gamma)},$$

then $\Phi(A; \alpha, \beta, \gamma) \leq \Phi(A; \alpha, \beta, \gamma)$ for any subsets α, β, γ of $\{1, 2, \ldots, n\}$.

Paper [28, 1964] deals with inequalities for principal subdeterminants of a special product of M-matrices. In this paper Ky Fan introduced the so-called Ky Fan product $A \odot B = \{ \begin{smallmatrix} a_{ii}b_{ii}, & i=j \\ -a_{ij}b_{ij}, & i \neq j \end{smallmatrix}$ and established that for two M-matrices A, B,

$$1 - \Phi(A \odot B; \alpha, \beta, \gamma) \leq (1 - \Phi(A; \alpha, \beta, \gamma))(1 - \Phi(B; \alpha, \beta, \gamma)).$$

For recent studies on some Ky Fan's results about M-matrices and the Ky Fan product see [66]. The importance of the Ky Fan product is that the set of M-matrices is closed under this product.

9. Ky Fan generalization of Szász's inequality

Suppose that $A \in \mathbb{M}_n$ is a positive-definite matrix and P_k $(1 \le k)$ is the product of all its $k \times k$ principal minors. The Szász inequality [64], as a generalization of Hadamard's determinantal inequality, says that $\prod_{i=1}^n a_{ii} = P_1 \ge \cdots \ge P_k^{1/\binom{n-1}{k-1}} \ge \cdots \ge P_n = \det(A)$. Ky Fan [31, 1967] strengthened the inequality for a certain class of matrices, named as GKK, including positive-definite matrices, totally positive matrices and M-matrices. By a GKK-matrix we mean one that all its principal minors are positive and the product of any two symmetrically situated almost principal minors is real and nonnegative. His result reads as follows.

Let $A \in \mathbb{M}_n$ be GKK and let $R_0 = I$, $R_k = Q_k^{1/\binom{p}{k}}$, $(1 \le k \le k)$, where $Q_k = \prod_{i_1<\cdots<i_k} A(\alpha_{i_1} \cup \cdots \cup \alpha_{i_k})$, in which α_i $(1 \le k \le p)$ are $p(\ge 2)$ pairwise disjoint possibly empty subsets of $\{1, \ldots, n\}$. Then $R_k^2 \ge R_{k-1}R_{k+1}$ and $Q_k^{1/\binom{p-1}{k-1}} \ge Q_{k+1}^{1/\binom{p-1}{k}}$ $(1 \le k \le p-1)$, which gives rise to the Szász inequality for $p = n$ and $\alpha_i = \{i\}$.

Ky Fan [41, 1992] generalized its former result by removing the assumption of the pairwise disjointness of the α_i, and showing that it remains true if $A - I$ is either positive-definite or an M-matrix.

10. Ky Fan extension of Kantorovich inequality

The Kantorovich inequality asserts that

$$\langle Hx, x\rangle\langle H^{-1}x, x\rangle \le (\lambda_1(H) + \lambda_n(H))^2/4\lambda_1(H)\lambda_n(H)$$

where x is a unit vector in $\mathbf{C}^n$ and H is an $n \times n$ positive-definite matrix. Ky Fan [29, 1966] generalized the inequality above by showing that if $0 < mI \le H \le MI$, $x_1, \ldots, x_m$ are vectors in $\mathbb{C}^n$ such that $\sum_{i=1}^n \|x_i\|^2 = 1$, then

$$\sum_{j=1}^m \langle H^p x_j, x_j\rangle \left[\sum_{j=1}^m \langle Hx_j, x_j\rangle\right]^{-p} \le (p-1)^{p-1}p^{-p}\left(b^p - a^p\right)^p((b-a)(ab^p - ba^p)^{p-1}\big)^{-1},$$

where p is any integer different from 0 and 1. In 1997 Mond and Pečarić [59] gave an operator version of Ky Fan's inequality above. Another extension of

Kantorovich inequality was given by Furuta [47]. Ky Fan proved that if H, K are positive operators on a Hilbert space, $H \geq K > 0$ and $MI \geq K \geq mI > 0$, then

$$\left(\frac{M}{m}\right)^{p-1} H^p \geq \frac{(p-1)^{p-1}}{p^p} \frac{(M^p - m^p)^p}{(M-m)(mM^p - Mm^p)^{p-1}} H^p \geq K^p$$

holds for all $p \geq 1$. The constant $\kappa_+(m, M, p) = \frac{(p-1)^{p-1}}{p^p} \frac{(M^p - m^p)^p}{(M-m)(mM^p - Mm^p)^{p-1}}$ is called the Ky Fan–Furuta constant in the literature; cf. [48].

11. Ky Fan minimax inequality

In his seminal work [32, 1972], Ky Fan presented the following result:

Let S be a nonempty compact convex subset of a Hausdorff topological vector space X, and let $f(x, y)$ be a function from $S \times S$ to the real numbers that is a lower semicontinuous function of y for each fixed $x \in S$ and is a quasi-concave function of x for each fixed $y \in S$. Then the following minimax inequality holds:

$$\min_{y \in S} \max_{x \in S} f(x, y) \leq \sup_{x \in S} f(x, x) \qquad \text{(Ky Fan minimax inequality)}.$$

It is equivalent to the Brouwer–Kakutani fixed point theorem and plays an essential role in several fields such as game theory, mathematical economics, variational inequalities, fixed point theory, control theory as well as nonlinear and convex analysis.

This inequality has evoked the interest of many mathematicians: Ferro [46] got a minimax inequality for vector-valued functions. Ding [16] obtained a minimax inequality by a generalized relatively Knaster–Kuratowski–Mazurkiewicz (R-KKM) mapping. Kuwano, Tanaka and Yamada [53] obtained a minimax inequality by using a unified scalarization. Also Hussain, Khan and Agarwal [51] used common fixed point theorem for a condensing map to obtain some Ky Fan type approximation theorems.

12. Sharpening of the von Neumann inequality

The well-known von Neumann inequality asserts that if A is a proper contraction (i.e, $\|A\| < 1$) acting on a Hilbert space, then $\|f(A)\| \leq 1$, where $f(A)$ is defined by applying the Riesz–Dunford functional calculus to a suitable analytic

function f. In several papers Ky Fan sharpened von Neumann's inequality by finding a bound of modulus less than 1 for $\|f(A)\|$; see [39, 1987].

(i) Ky Fan [35, 1979] proved the following inequality related to Julia's Lemma (for other results see [7, 1979]):

$$\left[I - \overline{f(z)}f(A)\right][I - f(A)^*f(A)]^{-1}[I - f(z)f(A)^*] \leq \frac{1-|f(z)|^2}{1-|z|^2}(I - \bar{z}A)(I - A^*A)^{-1}(I - zA^*)\,.$$

(ii) Ky Fan [38, 1987] established that

$$\|f(A)\| \leq [\|\mu_W(A)\| + |f(w)|]\,[1 + \|\mu_W(A)\|\,|f(w)|]^{-1},$$

in which $|w| < 1$ and μ_W is the Möbius transformation $\mu_W(z) = \frac{z-w}{1-\overline{w}z}$.
For recent develeopments concerning a multivariable von Neumann inequality see [49].

13. Operator Harnack inequalities

Ky Fan [40, 1988] provided two operator Harnack inequalities:
If $F(z)$ is a Hilbert space operator-valued analytic function on the open unit disk such that $F(0) = I$ and $\mathrm{Re}(F(z)) > 0$, then

$$(1-|z|)/(1+|z|)I \leq \mathrm{Re}(F(z)) \leq (1+|z|)/(1-|z|)I$$

and

$$-2|z|/(1-|z|^2)I \leq \mathrm{Im}(F(z)) \leq 2|z|/(1-|z|^2)I.$$

See also similar results of Ky Fan in [36, 1980].

Acknowledgement. The author would like to thank Dr. Farzad Dadipour and anonymous referees for carefully reading the manuscript and useful comments.

References

1. H. Alzer, *Converses of two inequalities of Ky Fan, O. Taussky and J. Todd*, J. Math. Anal. Appl. **161** (1991), 142-147.
2. H. Alzer, *The inequality of Ky Fan and related results*, Acta Appl. Math. **38** (1995), 305–354.
3. H. Alzer, *On an additive analogue of Ky Fan's inequality*, Indag. Math. (N.S.) **8** (1997), no. 1, 1–6.
4. H. Alzer, T. Ando and Y. Nakamura, *The inequalities of W. Sierpiński and Ky Fan*, J. Math. Anal. Appl. **149** (1990), no. 2, 497–512.
5. H. Alzer, S. Ruscheweyh and L. Salinas, *On Ky Fan-type inequalities*, Aequationes Math. **62** (2001), no. 3, 310–320.
6. A.R. Amir-Moéz and A. Horn, *Singular values of a matrix*, Amer. Math. Monthly **65** (1958), 742-748.
7. T. Ando and K. Fan, *Pick–Julia theorems for operators*, Math. Z. **168** (1979), no. 1, 23-34.
8. J.S. Aujla and F.C. Silva, *Weak majorization inequalities and convex functions*, Linear Algebra Appl. **369** (2003), 217-233.
9. E.F. Beckenbach and R. Bellman, *Inequalities*, Ergebnisse der Mathematik und ihrer Grenzgebiete, N. F., Bd. 30 Springer-Verlag, Berlin-Göttingen-Heidelberg 1961.
10. R. Bellman and K. Fan, *On systems of linear inequalities in Hermitian matrix variables*, in: V.L. Klee (Ed.), Convexity, The Proceedings of Symposia in Pure Mathematics, Vol. 7, American Mathematical Society, Providence, RI, 1963, pp. 1-11.
11. R. Bhatia, *Matrix Analysis*, Graduate Texts in Mathematics, 169. Springer-Verlag, New York, 1997.
12. P.S. Bullen, *A dictionary of inequalities*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 97. Longman, Harlow, 1998.
13. J.B. Chassan, *A statistical derivation of a pair of trigonometric inequalities*, Amer. Math. Monthly **62** (1955), 353-356.
14. C.-M. Cheng, R.A. Horn and C.-K. Li, *Inequalities and equalities for the Cartesian decomposition of complex matrices*, Linear Algebra Appl. **341** (2002), 219-237.
15. K.K. Chong, *Refinements of the Fan-Todd's inequalities*, Chinese Ann. Math. Ser. B **23** (2002), no. 1, 75-84.
16. X.P. Ding, *New generalized R-KKM type theorems in general topological spaces and applications*, Acta Math. Sin. (Engl. Ser.) **23** (2007), no. 10, 1869–1880.

17. J.L. Diaz-Barrero, J.J. Egozcue and J. Gibergans-Báguena, *On the Ky Fan inequality and some of its applications*, Comput. Math. Appl. **56** (2008), no. 9, 2279–2284.
18. S.S. Dragomir, *Some refinements of Ky Fan's inequality*, J. Math. Anal. Appl. **163** (1992), no. 2, 317–321.
19. K. Fan, *Sur les types homognes de dimensions* (French), C. R. Acad. Sci. Paris **211** (1940), 175-177.
20. K. Fan, *On a theorem of Weyl concerning eigenvalues of linear transformatioins I.*, Proc. Nat. Acad. Sci. U.S.A. **35** (1949), 652–655.
21. K. Fan, *On a theorem of Weyl concerning eigenvalues of linear transformations. II*, Proc. Nat. Acad. Sci. U. S. A. **36** (1950), 31-35.
22. K. Fan, *Maximum properties and inequalities for the eigenvalues of completely continuous operators*, Proc. Nat. Acad. Sci. U.S.A. **37** (1951), 760–766.
23. K. Fan, *Advanced Problems and Solutions: Solutions: 4430*, Amer. Math. Monthly **60** (1953), no. 1, 50.
24. K. Fan, *Inequalities for eigenvalues of Hermitian matrices*, Contributions to the solution of systems of linear equations and the determination of eigenvalues, pp. 131-139. National Bureau of Standards Applied Mathematics Series No. 39. U. S. Government Printing Office, Washington, D.C., 1954.
25. K. Fan, *A comparison theorem for eigenvalues of normal matrices*, Pacific J. Math. **5** (1955), 911-913.
26. K. Fan, *Some inequalities concerning positive-definite Hermitian matrices*, Proc. Cambridge Philos. Soc. **51** (1955), 414-421.
27. K. Fan, *Note on M-matrices*, Quart. J. Math. Oxford Ser. (2) **11** 1960, 43-49.
28. K. Fan, *Inequalities for M-matrices*, Nederl. Akad. Wetensch. Proc. Ser. A **67** (= Indag. Math. **26**) (1964), 602-610.
29. K. Fan, *Some matrix inequalities*, Abh. Math. Sem. Univ., Hamburg **29** (1966), 185-196.
30. K. Fan, *Inequalities for the sum of two matrices*, in Inequalities, O. Shisha, ed., Academic Press, New York, 1967.
31. K. Fan, *Subadditive functions on a distributive lattice and an extension of Szász's inequality*, J. Math. Anal. Appl. **18** (1967), 262-268.
32. K. Fan, *A minimax inequality and applications*, Inequalities, III (Proc. Third Sympos., Univ. California, Los Angeles, Calif., 1969; dedicated to the memory of Theodore S. Motzkin), pp. 103–113. Academic Press, New York, 1972.
33. K. Fan, *On real matrices with positive-definite symmetric component*, Linear and Multilinear Algebra **1** (1973), no. 1, 1-4.

34. K. Fan, *On strictly dissipative matrices*, Linear Algebra and Appl. **9** (1974), 223-241.
35. K. Fan, *Julia's lemma for operators*, Math. Ann. **239** (1979), no. 3, 241-245.
36. K. Fan, *Harnack's inequalities for operators*, General inequalities, 2 (Proc. Second Internat. Conf., Oberwolfach, 1978), pp. 333-339, Birkhäuser, Basel-Boston, Mass., 1980.
37. K. Fan, *Normalizable operators*, Linear Algebra Appl. **52/53** (1983), 253-263.
38. K. Fan, *Sharpened forms of an inequality of von Neumann*, Math. Z. **194** (1987), no. 1, 7-13.
39. K. Fan, *Applications and sharpened forms of an inequality of von Neumann*, Current trends in matrix theory (Auburn, Ala., 1986), 113-121, North-Holland, New York, 1987.
40. K. Fan, *Inequalities for proper contractions and strictly dissipative operators*, Linear Algebra Appl. **105** (1988), 237-248.
41. K. Fan, *Some inequalities for matrices A such that $A-I$ is positive-definite or an M-matrix*, Linear and Multilinear Algebra **32** (1992), no. 2, 89-92.
42. K. Fan and A.J. Hoffman, *Some metric inequalities in the space of matrices*, Proc. Amer.Math. Soc. **6** (1955) 111-116.
43. K. Fan and G.G. Lorentz, *An integral inequality*, Amer. Math. Monthly **61** (1954), 626-631.
44. K. Fan, O. Taussky and J. Todd, *Discrete analogs of inequalities of Wirtinger*, Monatsh. Math. **59** (1955), 73-90.
45. K. Fan and J. Todd, *A determinantal inequality*, J. London. Math. Soc. **30** (1955), 58-64.
46. F. Ferro, *A minimax theorem for vector-valued functions. II*, J. Optim. Theory Appl. **68** (1991), no. 1, 35-48.
47. T. Furuta, *Operator inequalities associated with Hölder-McCarthy and Kantorovich inequalities*, J. Inequal. Appl. **2** (1998), no. 2, 137–148.
48. T. Furuta, J. Mićić Hot, J.E. Pečarić and Y. Seo, *Mond–Pecaric Method in Ooperator Inequalities, Inequalities for Bounded Selfadjoint Operators on a Hilbert Space*, Monographs in Inequalities 1. Zagreb: Element, 2005.
49. A. Grinshpan, D. Kaliuzhnyi-Verbovetskyi, V. Vinnikov, and H.J. Woerdeman, *Classes of tuples of commuting contractions satisfying the multivariable von Neumann inequality*, J. Funct. Anal. **256**, 3035-3054.
50. R.A. Horn and C.R Johnson, *Topics in matrix analysis*, Corrected reprint of the 1991 original. Cambridge University Press, Cambridge, 1994.
51. N. Hussain A.R. Khan and R.P. Agarwal, *Krasnoselśkii and Ky Fan type fixed point theorems in ordered Banach spaces*, J. Nonlinear Convex Anal. **11** (2010), no. 3, 475-489.
52. M.V. Jovanović, T.K. Pogány and J. Sándor, *Notes on certain inequalities by Hölder, Lewent and Ky Fan*, J. Math. Inequal. **1** (2007), no. 1, 53–55.

53. I. Kuwano, T. Tanaka and S. Yamada, *Unified scalarization for sets and set-valued Ky Fan minimax inequality*, J. Nonlinear Convex Anal. **11** (2010), no. 3, 513–525.

54. N. Levinson, *Generalization of an inequality of Ky Fan*, J. Math. Anal. Appl. **8** (1964), 133–134.

55. B.-L. Lin, *Every waking moment Ky Fan (1914–2010)*, Notices Amer. Math. Soc. **57** (2010), no. 11, 1444-1447.

56. A.W. Marshall, I. Olkin and B.C. Arnold, *Inequalities: theory of majorization and its applications*, Second edition. Springer Series in Statistics. Springer, New York, 2011.

57. J.S. Matharu, M.S. Moslehian and J.S. Aujla, *Eigenvalue extensions of Bohr's inequality*, Linear Algebra Appl. **435** (2011), no. 2, 270–276.

58. D.S. Mitrinović, J.E. Pečarić and A.M. Fink,*Classical and New Inequalities in Analysis*, Mathematics and its Applications (East European Series), 61. Kluwer Academic Publishers Group, Dordrecht, 1993.

59. B. Mond and J.E. Pečarić, *Matrix inequalities for convex functions*, J. Math. Anal. Appl. **209** (1997), no. 1, 147-153.

60. M.S. Moslehian, R. Nakamoto and Y. Seo, *A Diaz-Metcalf type inequality for positive linear maps and its applications*, Electron. J. Linear Algebra **22** (2011), 179–190.

61. E. Neuman and J. Sandor, *On the Ky Fan inequality and related inequalities. II*, Bull. Austral. Math. Soc. **72** (2005), no. 1, 87–107.

62. A. Ostrowski, *Über die determinanten mit überwiegender Hauptdiagonale* (German), Comment. Math. Helv. **10** (1937), no. 1, 69-96.

63. J. Rooin, *An approach to Ky Fan type inequalities from binomial expansions*, Math. Inequal. Appl. **11** (2008), no. 4, 679–688.

64. O. Szász, *ber eine Verallgemeinerung des Hadamardschen Determinantensatzes* (German), Monatsh. Math. Phys. **28** (1917), no. 1, 253-257.

65. J. von Neumann, *Some matrix-inequalities and metrization of matrix-space*, Tomsk. Univ. Rev. 1, (1937), 286-300. [John von Neumann Collected Works (A. H. Taub, ed.), Vol. IV, pp. 205218. Pergamon, Oxford, 1962].

66. B.-Y. Wang, X. Zhang and F. Zhang, *On the Hadamard product of inverse M-matrices*, Linear Algebra Appl. **305** (2000), no. 1-3, 23-31.

67. H. Weyl, *Inequalities between the two kinds of eigenvalues of a linear transformation*, Proc. Nat. Acad. Sci. U. S. A. **35** (1949), 408-411.

68. X. Zhan, *Matrix Inequalities*, Lecture Notes in Mathematics, 1790. Springer-Verlag, Berlin, 2002.

69. F. Zhang, *Matrix Theory, Basic Results and Techniques*, Universitext, Springer-Verlag, New York, 1999.

Department of Pure Mathematics, Ferdowsi University of Mashhad, P.O. Box 1159, Mashhad 91775, Iran.

E-mail address: moslehian@ferdowsi.um.ac.ir, moslehian@member.ams.org

URL: http://profsite.um.ac.ir/~moslehian/